



5040 N. 15th Avenue, Suite 107 • Phoenix, AZ 85015
 Tel 602.759.1245 • Fax 602.759.8331 • www.CyrexLabs.com

ACCESSION #:
 TEST REQUISITION #:
 ACCOUNT #:
 NPI:
 DATE COLLECTED:
 DATE RECEIVED:
 DATE OF REPORT:
 SPECIMEN RECEIVED:

PRACTITIONER

PATIENT

NAME:
 DoB:
 ID:
 GENDER:

TEST	RESULTS				
	Normal	Equivocal*	Out of Range	Numeric Value	REFERENCE (ELISA Index)
ANTIBODY ARRAY 4					
Gluten-Associated Sensitivity & Cross-Reactive Foods					
Cow's Milk IgG+IgA Combined			X	2.33	0.4-2.0
Alpha-Casein & Beta-Casein IgG+IgA Comb.		X	.	1.81	0.2-2.3
Casomorphin IgG+IgA Combined	X			1.01	0.4-1.9
Milk Butyrophilin IgG+IgA Combined			X	2.84	0.3-1.3
American Cheese IgG+IgA Combined	X			1.42	0.4-2.2
Chocolate IgG+IgA Combined		X	.	2.18	0.3-2.4
Sesame IgG+IgA Combined	X			0.51	0.4-2.2
Hemp IgG+IgA Combined	X			1.03	0.6-1.8
Rye IgG+IgA Combined			X	2.68	0.5-2.1
Barley IgG+IgA Combined	X			1.09	0.4-1.8
Polish Wheat IgG+IgA Combined			X	2.73	0.4-2.1
Buckwheat IgG+IgA Combined			X	2.13	0.4-1.7
Sorghum IgG+IgA Combined	X			1.09	0.4-1.8
Millet IgG+IgA Combined	X			1.14	0.4-1.9
Spelt IgG+IgA Combined			X	4.51	0.2-2.0
Amaranth IgG+IgA Combined		X	.	1.84	0.3-2.5
Quinoa IgG+IgA Combined		X	.	1.46	0.3-1.7
Yeast IgG+IgA Combined		X	.	2.72	0.6-2.9
Tapioca IgG+IgA Combined	X			1.67	0.8-2.1
Oats IgG+IgA Combined			X	2.15	0.6-2.1
Coffee IgG+IgA Combined	X			0.75	0.5-2.5
Corn IgG+IgA Combined	X			1.58	0.6-2.6
Rice IgG+IgA Combined	X			1.60	0.6-2.4
Potato IgG+IgA Combined		X	.	1.62	0.7-2.0

*Reference ranges are calculated based on the mean \pm 2 standard deviations (SD). Results >1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.

Reviewed by: Dhiren Joshipura, M.D., Medical Director

Cyrex Laboratories is certified under the Clinical Laboratory Improvement Amendments of 1988 ("CLIA") as qualified to perform high-complexity clinical testing. Test result data on its own does not constitute a diagnosis. Only a physician or qualified healthcare professional should interpret the significance of a clinical lab test or make a diagnosis. This test was developed and its performance characteristics determined by Cyrex Laboratories, LLC. The names and titles of tests and arrays are for reference purposes only.